



Hyaluronic Acid Injection, Side Effects, and Complications

1. Introduction

Hyaluronic acid injection treatment results should be referred to the Chinese mock-up content of hyaluronic acid subdermal implant injection agent for treatment effect duration. The actual results may vary due to individual factors and can be checked during a follow-up visit 2 to 4 weeks after the injection.

2. Complications or side effects after treatment:

2.1 Skin condition after injection:

2.1.1 Minor bleeding may occur immediately after injection, which can be stopped by applying pressure with gauze for a few minutes. Avoid touching the injection site for the first 6 hours after injection.

2.1.2 Redness, bruising, itching, and other symptoms may occur within 1 to 2 days after injection. This is normal and will resolve on its own within a few days. Swelling in the lips may take longer to resolve, but this is not the final appearance result.

2.1.3 Avoid going to saunas or swimming pools within one week after injection.

2.1.4 If you had facial herpes before, recurrent herpes may occur at the injection site. This is normal and can be consulted or treated by your doctor.

2.2 Precautions after the treatment wound heals:

2.2.1 In rare cases, one or more injection sites may have an allergic reaction, which presents as persistent redness, itching, hard lumps, or bumps. If any of these symptoms occur, please return to the doctor for evaluation.

2.2.2 Delayed allergic reactions may occur in very few cases, presenting as redness and hard lumps months after injection. If this occurs, also return to the doctor for evaluation.