

















100.07.18 製定
102.04.11 第一次修訂
103.01.07 第二次修訂
藥衛 2

Antiplatelet Medication Information

Generic Name	Brand Name	Dosage form	Appearance
Aspirin	Bokey	100mg Capsules	
	Aspirin Protect	100mg Tablets	
	Ropal	100mg Tablets	
	Aspirin FC	100mg Tablets	
Ticlopidine	Licodin	100mg Tablets	
Clopidogrel	Plavix	75 mg Tablets	
	Clopidogrel	75mg Tablets	
Ticagrelor	Brilinta	60mg Tablets	
		90mg Tablets	



Generic Name	Brand Name	Dosage form	Appearance
Prasugrel	Efient	3.75mg Tablets	
		5mg Tablets	
Dipyridamole	Dipyridamole	25mg Tablets	
	Parotin	75mg Tablets	
	Persantin	75mg Tablets	

1. Why do I need this medicine?

Prevent platelet aggregation, reduce the risk of thrombosis, vascular obstruction, myocardial infarction, and stroke.

2. What information should I know about this medicine?

- (1) Do not use if you have ever been allergic to this drug or excipients.
- (2) If you have a history of peptic ulcer or using other anticoagulants, antiplatelet medications, anti-inflammatory painkillers, or have bleeding symptoms such as: bleeding gums, skin bruises, hematuria.
- (3) If you have renal or liver dysfunction, it may need dosage adjustment.
- (4) Check with your medical doctor before having any dental work or scheduling surgery.
- (5) Using this medicine while you are pregnant can harm your unborn baby. Use an



effective form of birth control to keep from getting pregnant during treatment.

- (6) Before surgical intervention, you should consult with your doctor to confirm the number of withdrawal days to reduce the risk of bleeding.

Brand Name	Precautions
Aspirin Protect Aspirin FC Bokey Ropal	Do not use in children or teenagers under the age of 20 to relieve symptoms of chickenpox or influenza. Patients with glucose-6-phosphate dehydrogenase deficiency should not use drugs containing salicylic acid.
Licodin	Please check the blood counts at the beginning of the drug and during the first three months.
Plavix Clopidogrel	Not recommended for use in children or adolescents under the age of 18.
Brilinta	Breastfeeding women should use it with caution, due to limited supporting evidence. Safety and effectiveness for use by children cannot be confirmed.

3. Possible Side Effects

Brand Name	Side Effects
Aspirin Protect Aspirin FC Bokey, Ropal	Gastrointestinal discomfort, diarrhea, allergies, skin rash, gastrointestinal bleeding, or bruising, dizziness, tinnitus, headache.
Licodin	Gastrointestinal upset, vomiting, loss of appetite, bleeding. Leukopenia may happen in some patients. Symptoms such as fever, chills, sore throat, and tiredness may appear at the initial stage.
Plavix Clopidogrel	Gastrointestinal discomfort, headache, dizziness, any abnormal bleeding or bruising on the body, difficulty stopping bleeding from wounds.
Brilinta	Difficulty breathing, headache, cough, dizziness, nausea, high blood pressure, diarrhea, fatigue, chest pain.
If the symptoms are severe and do not improve, please inform your doctor or pharmacist.	



4. What to do if you forget to take your medicine

- If you find that you have forgotten to take the medicine, please take it immediately, but if you think about it the next day, you only need to take the dose instructed by the doctor on the day, please do not take double dose, because it is easy to cause bleeding.
- If you forget to take the medicine for two or more days, please contact us immediately.

※Please consult your doctor or pharmacist if you have any questions.

Cheng Hsin General Hospital Medication Counseling Number

+886-2-2826-4400#5604#5605