

Cheng Hsin General Hospital

Angiotensin-converting-enzyme inhibitors (ACE inhibitors) Medication information

109.07.18 中文修訂 藥街 071

學名	英文商品名	劑量	禁衛 0/1
Amlodipine /Benazepril	Amtrel	5 mg/10 mg	Amtrel Tab 諾壓錠
Captopril	Ceporin	25 mg	CEPORIN 25 mg 血樂平錠劑
Fosinopril	Fonosil	10 mg	Fonosi I 10mg Tab 壓速利錠
Enalapril Maleate	Synbot	20 mg	Synbot 20mg Tab 欣保錠
Imidapril HCl	Tanatril	5 mg	TANATRIL 5 mg 田納滋錠劑
Ramipril	Tritace	2.5 mg	TRITACE 2.5 mg 心達舒錠劑
Ramipril	Tritace	10 mg	TRITACE 10mg Tab 心達舒錠10毫克

1. Why do I need this medicine?

Treatment for hypertension or heart failure

2. How is this drug best taken?

Use this drug as ordered by your doctor daily. Take Ceporin before meals with water.

3. What are some things I need to know or do while I take this drug?

- 1. To make sure the drug efficacy, do return to appointments for regular check-ups.
- 2. To prevent dizziness or headache, rise slowly if you have been sitting or lying down.
- 3. Avoid driving and doing other tasks or actions that call for you to be alert.
- 4. If you get pregnant while taking this drug, call your doctor right away. Do not breast-feed while you take this drug.
- 5. Manage high blood pressure with a healthy diet. Alcohol may interfere the effectiveness of ACEI, leading to dizziness and raising your risk of passing out.
- 6. If you are under diet with low sodium, drinking habits or other health conditions (especially diabetes mellitus, systemic lupus erythematosus, disease with liver or kidney), be sure to inform your doctor.

4. Possible Side Effect

- 1. If any of the following symptoms occur and cannot be relieved, seek medical attention. as soon as possible: dyspnea, facial or limbs swelling, , continuous dry cough, headache, fatigue, diarrhea or nausea. Take medicine with food or milk may relieve nausea.
- 2. If you fainted, stop medication and seek medical attention.

5. What do I do if I miss a dose?

- 1. Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- 2. Do not take double doses at the same time or extra doses.

6. How do I store this drug?

Store at room temperature in a dry place and keep out of the reach of children.

* Please consult your doctor or pharmacist if you have any questions.

Cheng Hsin General Hospital Medication Counseling Number

+886-2-2826-4400 #5604 #5605