



Oral Antiviral Treatment for Hepatitis B Drug Information

110.07.12 製訂
113.04 ENG 藥衛 099

Generic Name	Brand Name	Dosage form	Appearance
Entecavir	Baraclude	0.5mg Tablet	
		1mg Tablet	
Tenofovir disoproxil	Viread	300mg Tablet	
Tenofovir alafenamide	Vemlidy	25mg Tablet	

1. What is this drug used for?

To treat hepatitis B infection

2. How is this drug best taken?

Drug name	Medication time
Baraclude	Take on an empty stomach. (Take at least 2 hours before or 2 hours after a meal.) ◦
Viread	Take once daily at the same time of day. Take with or without food.
Vemlidy	

3. What are some things I need to know or do while I take this drug?

- (1) Discontinuation of medication may result in severe acute exacerbation of hepatitis B.
You cannot stop taking the drug on your own, and you must complete the course of

treatment according to the doctor's prescription.

- (2) After the treatment, regular follow-up visits should still be made for clinical, blood and imaging tests.
- (3) Please inform your doctor if you are under one of following condition
 - a) Taking any other medications or health supplements
 - b) Pregnant, plan to become pregnant or breastfeeding
 - c) Kidney disease, liver disease, or ever had liver transplants

4. Possible Side Effect

Fatigue, headache, diarrhea, nausea, bloating, indigestion, gastrointestinal discomfort

5. What do I do if I miss a dose?

- (1) Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- (2) Do not take double doses at the same time or extra doses.

6. How do I store this drug?

Please store in a dry and cool place below 30°C.

*This leaflet cannot provide complete drug information.
Please consult doctors or pharmacist if you have any questions.

Cheng Hsin General Hospital Medication Counseling Number
+886-2-2826-4400 #5604 #5605