

# **Cheng Hsin General Hospital**

# **Angiotensin II Receptor Blockers Drug Information**

109.11.19 修訂 113.05 二修 ENG 藥衛 075

### Single Ingredient

Generic Name	Brand Name	Dosage form	Appearance
Candesartan	Blopress	8 mg	博脈舒錠劑 Blopress 8mg Tab
Valsartan	Diovan	160 mg	名 将安穏膜衣紋160毫克 Diovan FC 160mg Tab
Azilsartan Medoxomil	Edarbi	40 mg	易得平錠 Edarbi 40mg Tab
Olmesartan	Olmetec	40 mg	雅縣膜衣錠 OLMETEC 40mg Tab

# Multi-Ingredient

Amlodipine/ Valsartan	Exforge	5/80 mg	易安穩縣衣紋 Exforge 5/80mg Tab
Losartan/ Hydrochlorothiazide	Hyzaar	100/12.5 mg	好悅悠默衣錠 llyzaar Tab
Amlodipine/ Olmesartan	Sevikar	5/40mg	行脈康膜衣紋 SEVIKAR 5/40mg Tab
Candesartan/ Hydrochlorothiazide	Blopress Plus	16/12.5 mg	機脈舒加強錠剤 BLOPRESS PLUS Tab

# 1. Why do I need this medicine?

To treat high blood pressure or heart failure. Combination drugs contains diuretics (Hydrochlorothiazide) or calcium channel blockers(Amlodipine), with synergistic

antihypertensive effect.

#### 2. How is this drug best taken?

Please take it with water and follow the doctor's instruction. Do not start, stop, or change the dose of any drug without checking with your doctor.

#### 3. What are some things I need to know or do while I take this drug?

- (1) Make sure if you are allergic to this drug before taking it. If you have signs of an allergic reaction get medical help immediately.
- (2) Tell your doctor about all of your drugs (including diuretics, lithium, potassium supplements, or other blood pressure-lowering medications) to prevent drug-drug interactions.
- (3) Tell your doctor about all of your health problems, including kidney, liver, heart disease or diabetes.
- (4) Tell your doctor in advance if you are pregnant, planning to get pregnant or breastfeeding.

#### 4. Possible Side Effect

Common: dizziness, postural dizziness, hypotension, postural hypotension

Rare: Hyperkalemia, headache, syncope, cough, diarrhea, renal insufficiency, allergic reaction, angioedema

#### 5. What do I do if I miss a dose?

- (1) Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- (2) Do not take double doses at the same time or extra doses.

## 6. How do I store this drug?

Store at room temperature.

\*This leaflet cannot provide complete drug information. Please consult doctors or pharmacist if you have any questions.

Cheng Hsin General Hospital Medication Counseling Number +886-2-2826-4400 #5604 #5605