

Cheng Hsin General Hospital

Vaginal Suppository Drug Information

109.07.18 中文修訂 113.02 ENG 藥衛 048

	D 131		113.02 ENO 崇稱 040
Generic Name	Brand Name	Dosage Form	Appearance
Clotrimazole	Clomazole	500mg Vag Tab	Clomazole 500mg Vag Tab Clomazole 500mg Vag Tab Company of the
Fenticonazole	Lomexin	200mg Vag Cap	各微欣陰道軟膠囊 Lomexin 200mg Vag Tab
Sertaconazole	Zalain	500mg Vag Tab	達來陰道錠 Zalain 500mg Vag Tab Salain 500mg Vag Tab Zalain 500mg Vag Tab
Estriol	Esvatin	0.5mg Vag Tab	被詩婷陰道錠 Esvatin Vag 0.5mg Tab

1. Why do I need this medicine?

- (1) The main components of Clomazole, Lomexin and Zalain vaginal suppositories are antifungal agents. They are used to treat vaginal infections caused by fungi, predominantly Candida albicans, commonly known as yeast infections.
- (2) The main component of Esvatin vaginal tablet is estriol. It is used to treat vaginitis, cervicitis, and cervical-vaginal erosion.

2. How is this drug best taken?

- (1) It is recommended to use vaginal suppositories after bathing before bedtime.
- (2) Wash your hands thoroughly and dry them with a clean tissue.

- (3) Tear open the external packaging of the vaginal suppository and moisten it with warm water.
- (4) Following the prescribed dosage, squat or lie down with your knees bent, then use your finger to insert the suppository into the vagina to a depth of about one finger.
- (5) Wash your hands thoroughly again after use.

3. What are some things I need to know or do while I take this drug?

- (1) Vaginal suppositories should not be used on other parts of the body and should not be swallowed.
- (2) Complete the entire course of treatment and do not stop the medication arbitrarily.
- (3) Continue using it even during menstruation.

4. Possible side effects

If you experience persistent irritation, such as burning or itching, please inform your doctor immediately for evaluation.

5. What do I do if I miss a dose?

Take it as soon as you remember. If it's close to your next dose, skip the missed one and continue with your regular schedule. Do not use 2 doses at the same time and be sure to inform your doctor.

6. How do I store this drug?

Store in a cool and dry place at room temperature, away from direct sunlight. Keep out of reach of children.

*This leaflet cannot provide complete drug information. Please consult a doctor or pharmacist if you have any questions.

Cheng Hsin General Hospital Medication Counseling Number +886-2-2826-4400 #5604 #5605