




Evac Enema Drug Information

109.07.18 中文修訂
113.02 ENG 藥衛 062

Generic Name	Brand Name	Dosage Form	Appearance
Monosodium phosphate, Disodium phosphate	Evac	118 mL Enema (139.1mg/31.8mg per mL)	

1. Why do I need this medicine?

This enema solution has a laxative effect, allowing for the safe and direct clearance of the colon contents within 2 to 15 minutes. It is suitable for relieving constipation, emptying the colon contents before colonoscopy, and preparing the bowel before abdominal surgery.


2. How is this drug best taken?

Evac enema should be used as directed by your doctor. Adults should use one bottle per dose, while children aged 2 to 12 should use half the dose.


Steps for using Evac enema:

※ You may apply petroleum jelly to the soft tube before gently inserting the tube into the rectum. If there is resistance while squeezing the bottle to inject the solution, stop immediately to prevent damage to the intestinal wall.


1. Remove the bottle cap.



2. Screw the tube onto the bottle.



3. With one hand holding the direction of the tube steady, use the other hand to gently insert it into the rectum until it reaches the bottle's neck.



4. Squeeze from the bottom of the bottle towards the front until the solution is completely injected.
5. Hold the thumb steady and pull the soft tube out directly. Discard the enema kit properly.



3. What are some things I need to know or do while I take this drug?

- (1) Do not use any laxatives or Evac enema if you experience abdominal pain, nausea, or vomiting.
- (2) Frequent use of this medication may lead to dependency.
- (3) This medication is not recommended for children under two years old.
- (4) Use with caution if you have impaired kidney function, heart disease, electrolyte imbalance, or are currently taking calcium channel blockers or diuretics.

4. Possible side effects

This medication may cause hypocalcemia, hyperphosphatemia, hypernatremia, or acidosis. If you experience discomfort or gastrointestinal bleeding after use, please stop using it immediately and seek medical attention.

5. What do I do if I miss a dose?

Make sure you take the medication as prescribed at the correct times to facilitate examinations or surgeries. Never take double the dose at once. If you forget to take it at the designated time, please consult your doctor.

6. How do I store this drug?

Please store at room temperature in a dry place. Keep out of reach of children.

*This leaflet cannot provide complete drug information.
Please consult a doctor or pharmacist if you have any questions.

Cheng Hsin General Hospital Medication Counseling Number
+886-2-2826-4400 #5604 #5605