









101.10.19 製定  
102.04.11 第一次修訂  
104.03.02 第二次修訂  
104.12.24 第三次修訂  
藥衛 47

# Rectal Suppository Medication Information

Generic Name	Brand Name	Dosage	Indication	Appearance
Na oleate, Polidocanol, Chlorcarvachol	Alcos-Anal Supp	200 mg, 20 mg, 0.7 mg	hemorrhoids	
Mesalazine	Asacol Supp	500 mg	Ulcerative Colitis, Ulcerative proctitis, Proctosigmoiditis	
Bisacodyl	Bisacodyl Supp	10 mg	constipation	
Domperidone	Domper Supp	10 mg	nausea and vomiting	
Polycresol -sulfonate, Cinchocaine	Faktu Supp	100 mg, 2.5 mg	hemorrhoids	
Diclofenac Na	Voren Supp	12.5 mg	antipyretic analgesic	

## 1. How to use this medicine?

- (1) Wash your hands and dry thoroughly
- (2) Removing plastic wrap from the suppository. Moisturize the end of the suppository with water if necessary.
- (3) Lie on one side with one leg bent and the other straightened.
- (4) Gently and firmly push suppository into your rectum for about 5 centimeter (one to two knuckles deep) to prevent slipping out.
- (5) Lie with your legs closed for a few minutes.
- (6) Wash your hands right away with soap and warm water.

## 2. What information should I know about this medicine?

- (1) Follow the prescription and instruction by the doctor precisely.



101.10.19 製定  
102.04.11 第一次修訂  
104.03.02 第二次修訂  
104.12.24 第三次修訂  
藥衛 47

- (2) Keep out of the reach of children.
- (3) Rectal suppositories cannot be used to other routes.

#### **4. Possible Side Effects**

Inform your doctor when you have rectal burning or irritation.

#### **5. What should I do if I forget to take my medication?**

- (1) If you found that you forget to take your medication, please take the medication as soon as you remember.
- (2) If it is already close to the time for next dose, please skip this one dose, and take the next dose of medication on time.
- (3) Remember not to take double dosing in a single medication intake.

#### **6. Storage and Stability**

Store the suppositories in a cool place to prevent melting.

※ Please consult your doctor or pharmacist if you have any questions.

Cheng Hsin General Hospital Medication Counseling Number

+886-2-2826-4400 #5604 #5605