




Nincort Oral Gel Drug Information

109.07.18 中文修訂
113.02 ENG 藥衛 019

Generic Name	Brand Name	Dosage Form	Appearance
Triamcinolone acetonide	Nincort	0.1% 6 gm Oral Gel	

1. Why do I need this medicine?

Treatment for inflammatory or ulcerative oral lesions.

2. How is this drug best taken?

Apply once daily to the affected area, preferably before bedtime. Apply 2-3 times daily if symptoms are severe. Avoid eating or drinking immediately after application.

3. What are some things I need to know or do while I take this drug?

- (1) Stop using this medicine if there is purulent infection locally.
- (2) Stop using this medicine immediately if allergic reactions occur.
- (3) If there is no significant improvement at the affected area after seven days of use, seek medical examination.
- (4) Patients with tuberculosis, diabetes, or gastric ulcers should not use this medication.

4. Possible side effects

Rare.

5. What do I do if I miss a dose?

Take it as soon as you remember. If it's close to your next dose, skip the missed one and continue with your regular schedule. Do not use 2 doses at the same time and be sure to inform your doctor.

6. How do I store this drug?

Store in a dry place at a temperature ranging from 15-25 °C. Do not use if the product has expired or shows any signs of deterioration.

*This leaflet cannot provide complete drug information.
Please consult a doctor or pharmacist if you have any questions.

Cheng Hsin General Hospital Medication Counseling Number
+886-2-2826-4400 #5604 #5605