




Premarin Vag Cream Drug Information

109.07.18 中文修訂
113.02 ENG 藥衛 032

Generic Name	Brand Name	Dosage Form	Appearance
Conjugated estrogen	Premarin	14gm Vag Cream	

1. Why do I need this medicine?

Treatment for atrophic vaginitis and kraurosis vulvae due to menopause.

2. How is this drug best taken?

- (1) Remove the cap from the tube.
- (2) Screw the applicator onto the tube.
- (3) Gently squeeze the tube from the bottom to force the prescribed amount of medication into the applicator, following the markings on it.
- (4) Unscrew the applicator from the tube.
- (5) Lie down with knees drawn up. Gently insert the applicator deeply into the vagina. Press the plunger to release the medication completely into the vagina.

3. What are some things I need to know or do while I take this drug?

Pull the plunger to remove it from the barrel and then wash with mild soap and warm water after use. Do not boil or use hot water.

4. Possible side effects

Nausea, vomiting, abdominal cramps, headache, changes in menstrual periods, swelling, changes in weight, and other symptoms.

5. What do I do if I miss a dose?

Take it as soon as you remember. If it's close to your next dose, skip the missed one and continue with your regular schedule. Do not use 2 doses at the same time and be sure to inform your doctor.

6. How do I store this drug?

The recommended storage temperature is 20~25°C. Temperature variations between 15~30°C are acceptable for short-term travel.

*This leaflet cannot provide complete drug information.
Please consult a doctor or pharmacist if you have any questions.

Cheng Hsin General Hospital Medication Counseling Number
+886-2-2826-4400 #5604 #5605