




## Rivast Oral Solution Drug Information









110.07.12 修訂  
113.04 ENG 藥衛 096





Generic Name	Brand Name	Dosage form	Appearance
Rivastigmine	Rivast Oral Solution	240mg/120ml	

### 1. What is this drug used for?

To treat dementia in people with Alzheimer's disease or dementia in people with Parkinson's disease.

### 2. How is this drug best taken?

 <p>① Check the packaging set (one each for potion, dosing syringe, base and plugged straw).</p>	 <p>② Press the cap and turn it anticlockwise</p>	 <p>③ Insert the plugged straw into the bottle neck. Make the top fit tightly with the mouth of the bottle.</p>	 <p>④ Take the syringe and put it in the adaptor opening</p>
 <p>⑤ Fill the syringe with a small amount of solution by pulling the piston out</p>	 <p>⑥ Turn the syringe upside down, pull the piston downward to remove gas.</p>	 <p>⑦ Push the piston upward in order to remove any possible bubbles.</p>	 <p>⑧ Slowly drain the excess dose back into the bottle.</p>

 <p>⑨Align the scale line on the push rod with the barrel body to determine the required amount of medicine.</p>	 <p>⑩Peel off the double-sided tape on the side of the base.</p>	 <p>⑪Glue the base to the bottom of the potion bottle.</p>	 <p>⑫After use, cap the bottle and insert the dosing syringe back into the base.</p>
---	---	--	---

### 3. What are some things I need to know or do while I take this drug?

- (1) Patients with heart disease, asthma, or obstructive pulmonary disease need to use it with caution.
- (2) This medicine may increase gastric acid secretion and may worsen urinary tract obstruction or seizures.

### 4. Possible Side Effect

- (1) Common adverse reactions include nausea, vomiting, diarrhea, loss of appetite, weight loss, and confusion.
- (2) Gastrointestinal adverse reactions tend to occur during dose adjustment. If discomfort persists, you can adjust to the previous dosage first, and tell your doctor.

### 5. What do I do if I miss a dose?

- (1) Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- (2) Do not take double doses at the same time or extra doses.

### 6. How do I store this drug?

Store at room temperature under 25 °C. Do not refrigerate and keep it from heat.

\*This leaflet cannot provide complete drug information.  
Please consult doctors or pharmacist if you have any questions.

Cheng Hsin General Hospital Medication Counseling Number  
+886-2-2826-4400 #5604 #5605