




Sutrol Vag Gel Drug Information

109.07.18 中文修訂
113.02 ENG 藥衛 038

Generic Name	Brand Name	Dosage Form	Appearance
Metronidazole	Sutrol	0.75% 25gm Vag Gel	

1. Why do I need this medicine?

Treatment for vaginal infections such as vaginal trichomoniasis and abnormal vaginal discharge.

2. How is this drug best taken?

- (1) Screw the applicator onto the tube.
- (2) Squeeze the medication into the applicator until it is full.
- (3) While sitting or lying down, hold the outer part of the applicator, gently insert it deep into the vagina and deliver the medication.
- (4) Clean the applicator with warm water after use. Do not boil or use hot water for cleaning.

3. What are some things I need to know or do while I take this drug?

- (1) Patients who are prone to edema or are taking corticosteroids should take special care when using this medication.
- (2) Patients with a history of hematologic disorders should have their white blood cell count measured before and after treatment.
- (3) Avoid using aluminum products, such as needles or syringes, in contact with the medication.
- (4) If retreatment is necessary, there should be a 4 to 6-week interval between treatment courses.
- (5) For patients who have genital infections, their sexual partners should also receive treatment.

4. Possible side effects

The skin may experience temporary redness, mild dryness, a burning sensation, and irritation after use.

5. What do I do if I miss a dose?

Take it as soon as you remember. If it's close to your next dose, skip the missed one and continue with your regular schedule. Do not use 2 doses at the same time and be sure to inform your doctor.

6. How do I store this drug?

Store at a room temperature below 25° C and keep it away from light. Use it before the expiration date.

*This leaflet cannot provide complete drug information.
Please consult a doctor or pharmacist if you have any questions.

Cheng Hsin General Hospital Medication Counseling Number
+886-2-2826-4400 #5604 #5605